

*ins
A1*
*cls
E'*

Title: Prevention and treatment of Alzheimer's disease

Field of the invention

The present invention relates to prevention and treatment of Alzheimer's disease (AD). More specifically, the invention relates to use of a non-wild type protofibril or compound(s) with protofibril forming ability for active immunisation in the purpose of treating or preventing AD. The invention further relates to a peptide, A β -Arc, with high protofibril forming activity as well as several applications thereof, such as antibodies against said peptide for passive immunisation against AD.

Background of the invention

Alzheimer's disease (AD) is a progressive disease known generally as senile dementia. The disease falls into two categories, namely late onset and early onset. One form of this latter AD type runs in families and it is known as familial AD.

Both types of AD are characterized by two types of lesions in the brain: senile plaques and neurofibrillary tangles. Senile plaques are areas of disorganized neuropil up to 150 mm across with extracellular amyloid deposits at the center. Neurofibrillary tangles are intracellular deposits consisting of two filaments twisted about each other in pairs.

*sub
cl*

A β also referred to as amyloid β peptide (A β P) is a highly aggregating small polypeptide having a molecular weight of approximately 4,500. This protein is a cleavage product of a much larger precursor protein referred to as amyloid precursor protein (APP). The A β protein comprises 39 - 42 amino acids. There are at least five distinct isoforms of APP: 563, 695, 714, 751, and 770 amino acids, respectively (Wirak et al. (1991)). The A β protein segment comprises approximately half of the transmembrane domain and approximately the first 28 amino acids of the extracellular domain of an APP isoform.

APP is a transmembrane protein which is highly expressed in all parts of the body, and which has several important biological functions. Proteolytic processing of APP in vivo is a normal physiological process. Carboxy-terminal truncated forms of APP695, APP751, and APP770 are present in brain and cerebrospinal fluid (Palmer et al. (1989)) (Weidemann et al (1989)). There are probably two main metabolic pathways: one non-amyloid-forming and one amyloid-forming pathway. The amyloid forming non-normal pathway produces the A β protein polypeptide which is prone to form dense amyloidogenic aggregates that

are resistant to proteolytic degradation and removal. The resultant A β protein aggregates presumably are involved in the formation of the abundant amyloid plaques and cerebrovascular amyloid that are the neuropathological hallmarks of AD.

In AD brains, the A β peptide forms virtually insoluble amyloid fibrils that accumulate into senile plaques. The A β fibrillization process is a complex multistep reaction. A group of distinct intermediary A β species of the fibrillization reaction, the protofibrils, were recently identified (Walsh et al. (1997)), (Walsh et al. (1999)), (Harper et al. (1999)).

The most common A β form in cerebrospinal fluid (CSF) and plasma comprises 40 amino acids (A β 40), but an A β comprising 42 amino acids (A β 42) is the most common form in plaques (Scheuner et al. (1996)). This longer form tends to aggregate more rapidly and it is believed that it is more pathogenic than A β 40.

Many patients get Alzheimer's disease spontaneously with unknown etiology, but there are also several hereditary components involved. Disease-causing mutations in genes on chromosomes 1, 14, and 21, respectively, have been discovered, and these mutations might explain as much as 50% of disease forms starting very early (<50 years) (St. George-Hyslop et al. (1987)), (Sherrington et al. (1995)).

The first gene associated with Alzheimer's disease was the gene encoding the amyloid precursor protein APP on chromosome 21. Different mutations of this gene result in unusual hereditary forms of the disease. Several pathogenic mutations have been identified in the (APP) gene, all located close to the major APP processing sites. These processing sites are either located adjacent to the boundaries of the A β domain in APP (the β - and γ -secretase sites) or within the A β sequence itself (α -secretase site).

The only known AD mutation close to the β -secretase site, the Swedish mutation (Mullan, et al., (1992)), discloses a double mutation (Lys670Asn/Met671Leu) of the APP gene in a large Swedish family, in which family the disease starts early and has a high penetrating power. The mutation produces a large increase of A β production, an elevation of both A β 42 and A β 40 in plasma from mutation carriers and in conditioned cell media.

Other APP mutations have been described. All result in Alzheimer's disease with an early age of onset having an autosomal dominant heredity pattern. Pathogenic mutations within the A β sequence, located close to the α -secretase site, result in a phenotype different from AD, with massive amyloid accumulation in cerebral blood vessel walls. Two mutations at codons 692 and 693, namely the Dutch (Glu693Gln) and the Flemish (Ala692Gly) mutations, have been reported (Levy et al. (1990)), (van Broeckhoven et al. (1990)), (Hendriks et al. (1992)). Patients having these mutations suffer from cerebral haemorrhage and vascular symptoms. The vascular symptoms are caused by aggregation of A β in blood vessel walls (amyloid angiopathy). A third pathogenic intra-A β mutation was recently discovered in an Italian family (E693K), with clinical findings similar to the Dutch patients (Tagliavini, et al. (1999)).

Different pathogenic mechanisms have been proposed for the Dutch and Flemish mutations. It has been observed that the Flemish mutation leads to increased A β levels while a reduced ratio of A β 42/40 was seen in media from cells transfected with the Dutch mutation (De Jonghe, et al.(1998)). Investigations of synthetic A β peptides have indicated that the Dutch mutation, but not the Flemish, accelerates the fibril formation compared to wild-type (wt) peptide (Walsh et al. (1997)).

As reported by Kamino et al. 1992, another APP E693 variant wherein Glu is substituted for Gly at APP E693, has previously been seen in one individual. It could not be unambiguously determined to be responsible for AD, though. This case originated from a family with similar clinical characteristics for AD and definitive AD was confirmed at autopsy. However, in this family the mutation could only be detected in one of two demented siblings.

Mice transgenic for APP mutations show many of the pathological features of Alzheimer disease, including deposition of extracellular amyloid plaques, astrogliosis and neuritic dystrophy. In recent studies by (Schenk et al (1999)) it was reported that immunization with A β 42 wild-type peptide is both preventive in transgenic mice, but also that A β containing plaques can be greatly reduced in the brain of transgenic mice immunized with the peptide.

However, due to the large costs and suffering that are associated with Alzheimer's disease, there is still a need for improved methods for treatment and prevention thereof.

Likewise, there is a need for a method for screening compounds that could constitute a part of future pharmaceutical preparations for treating and perhaps curing Alzheimer's disease.

Summary of the invention

The present invention relates to an active immunisation against AD which will have a much more profound effect in the treatment of Alzheimer's disease, than using the wild-type peptide. Immunization according to the invention will yield antibodies directed to protofibrils, as the immunogen is a protofibril or compound(s) with greatly increased protofibril formation properties. These antibodies, generated in the periphery, will cross the blood brain barrier and mediate clearance of A β in the brain in a protofibril state.

In present invention use is made of a pathogenic AD mutation at codon 693 (Glu693Gly), named the 'Arctic mutation', located within the A β peptide domain of the APP gene, more closely position 22 of the A β -Arc peptide. Carriers of this mutation develop progressive dementia with clinical features typical of AD without symptoms of cerebrovascular disease. Said AD is distinctly characterised by accelerated formation of protofibrils comprising mutated A β peptides (40Arc and/or 42Arc) compared to protofibril formation of wild type A β peptides.

Thus, in a first aspect the invention relates to use of a non-wild type protofibril or compound(s) with protofibril forming ability for immunisation for prevention or treatment of Alzheimer's disease (AD). Preferably, these protofibril or compound(s) have enhanced protofibril forming ability and/or enhanced immunogenicity compared to the wild-type counterparts. Protofibril chemistry has been described by, inter alia, Serpell (2000).

Preferably, the protofibril or compound(s) with protofibril forming ability comprises the following amino acid sequence KLVFFAEDV. The A β 1-42 fibrillisation process involves transitional conformation changes from α -helix via random coil to β -sheet. The stable α -helix sequence of residues 16-24 (KLVFFAEDV) apparently plays an important role in this process.

The protofibril or compound(s) with protofibril forming ability may be mutated or modified in relation to corresponding wild-type counterparts. Changes in the KLVFFAEDV

SECTION 2

sub
cg

sequence will affect the fibrillisation process. For example, changes of the charged amino acids Glu22 and Asp23 into neutral amino acids will induce a random coil structure in the A β peptide. Furthermore, deprotonation of other amino acids such as Asp7, Glu11 and His 6, 13 and 14 in the N-terminal end, has been suggested to destabilize the α -helix, leading to initiation of the fibrillation process. Another example is mutations leading to increased immunogenicity in man by using amino acids from mouse A β at specific positions, e.g. Gly 5, Phe10, Arg13. Furthermore, amino acid 13 in A β is known to be part of a heparan sulphate binding motif (13-16; His, His, Gln, Lys) in human, which has been speculated to be involved in AD disease mechanism (inflammation) (Giulian et al. (1998)). In mouse, His 16 is exchanged for Arg 13 destroying the heparan sulphate binding site. Interestingly, mice have never been observed to develop AD. Hence, the use of A β -Arc/Arg13 as an immunogen would be a way to lower possible inflammatory side effects, elicited with A β peptides with intact heparan sulphate binding motif.

Preferably, the protofibril or compound(s) with protofibril forming ability comprises an A β peptide (β -amyloid protein) and repeats thereof, such as dimeric, oligomeric or multimeric forms). In a preferred embodiment the protofibril or compound(s) with protofibril forming ability comprises a A β peptide related to AD. In another embodiment the protofibril or compound(s) with protofibril forming ability comprises α -synuclein.

There exists a form of dementia characterised by patients having clusters in the brain of a structure called Lewy bodies. This form of dementia comprises about 20% of all dementia. Patients with Lewy bodies show, inter alia, Parkinson symptoms with progressive cognitive dysfunction. However, some patients also exhibit Alzheimer symptoms and this is called "Lewy variant of Alzheimer". The main component of the Lewy bodies is the protein α -synuclein. Two mutations in α -synuclein have been identified Ala53Thr and Ala30Pro. These mutations lead to dominant heritage of Parkinson's disease. These mutations affect the structure/solubility of α -synuclein and leads to formation of protofibrils. (Conway et al. (2000)).

The A β peptide is preferably A β -Arc as disclosed in SEQ ID NO 1. A β -Arc comprises 39, 40 or 42 amino acids but may also be shorter as long as the protofibril forming ability is maintained.

The profibril or compound(s) with protofibril forming ability may be used in combination with A β peptides having known mutations, such as the Dutch, Flemish, Italian mutation described above as well as the Iowa mutation (D694N) (Grabowski et al., 2001).

The A β peptide may comprise one or more of these and/or other mutations. Alternatively, a cocktail of different A β peptides with different mutations is used.

In a second aspect, the invention relates to a peptide, A β -Arc, having the amino acid sequence disclosed in SEQ ID NO 1 comprising a glycine at position 22 instead of glutamic acid compared to wild type A β peptide. The peptide may be natural, synthetic or recombinantly produced. For the purposes of the invention the peptide may be used in monomeric, dimeric, oligomeric, protofibril or multimeric form.

The invention also relates to nucleic acid encoding the above peptide as well as a vector comprising the nucleic acid. The vectors for expressing the polypeptides of the invention require that the nucleic acid be "operatively linked." A nucleic acid is operatively linked when it is placed into a functional relationship with another nucleic acid sequence.

This vector may be inserted in a host cell. Such a host cell can be used to recombinantly produce the peptide of the invention for pharmaceutical or diagnostic use as well for research purposes. The peptide may also be produced synthetically and be purified by HPLC, RP-HPLC, SEC-HPLC.

In a further aspect, the invention relates to a transgenic non-human animal comprising the above vector. Furthermore, the invention relates to a transgenic non-human animal comprising a vector comprising the entire APP gene corresponding to NCBI database, accession no XM_009710, Homo sapiens amyloid β (A4) precursor protein (protease nexin-II, Alzheimer's disease)(APP), mRNA. However, the APP gene for use in the invention comprises the Arctic mutation, i.e. nucleotide number 2225 is mutated from A to G leading to an amino acid substitution from Glutamic acid to Glycine. The transgenic animal may be used for modelling Alzheimer's disease and testing for therapeutic treatment efficacy. This transgenic animal will bear the entire APP gene comprising the Arctic mutation. This gene is preferably under control of a strong promoter, such as the prion-promoter. The APP gene may contain further mutations, besides the Arctic mutation.

Time	Lat	Long	Alt	Wind	Temp	Hum	Clouds	Vis	Remarks
0000	10° 00' N	155° 00' E	1000	000	25.0	80	000	10	Clear
0100	10° 05' N	155° 05' E	1000	000	25.0	80	000	10	Clear
0200	10° 10' N	155° 10' E	1000	000	25.0	80	000	10	Clear
0300	10° 15' N	155° 15' E	1000	000	25.0	80	000	10	Clear
0400	10° 20' N	155° 20' E	1000	000	25.0	80	000	10	Clear
0500	10° 25' N	155° 25' E	1000	000	25.0	80	000	10	Clear
0600	10° 30' N	155° 30' E	1000	000	25.0	80	000	10	Clear
0700	10° 35' N	155° 35' E	1000	000	25.0	80	000	10	Clear
0800	10° 40' N	155° 40' E	1000	000	25.0	80	000	10	Clear
0900	10° 45' N	155° 45' E	1000	000	25.0	80	000	10	Clear
1000	10° 50' N	155° 50' E	1000	000	25.0	80	000	10	Clear
1100	10° 55' N	155° 55' E	1000	000	25.0	80	000	10	Clear
1200	11° 00' N	156° 00' E	1000	000	25.0	80	000	10	Clear
1300	11° 05' N	156° 05' E	1000	000	25.0	80	000	10	Clear
1400	11° 10' N	156° 10' E	1000	000	25.0	80	000	10	Clear
1500	11° 15' N	156° 15' E	1000	000	25.0	80	000	10	Clear
1600	11° 20' N	156° 20' E	1000	000	25.0	80	000	10	Clear
1700	11° 25' N	156° 25' E	1000	000	25.0	80	000	10	Clear
1800	11° 30' N	156° 30' E	1000	000	25.0	80	000	10	Clear
1900	11° 35' N	156° 35' E	1000	000	25.0	80	000	10	Clear
2000	11° 40' N	156° 40' E	1000	000	25.0	80	000	10	Clear
2100	11° 45' N	156° 45' E	1000	000	25.0	80	000	10	Clear
2200	11° 50' N	156° 50' E	1000	000	25.0	80	000	10	Clear
2300	11° 55' N	156° 55' E	1000	000	25.0	80	000	10	Clear

36

Another aspect of the invention relates to a pharmaceutical composition, comprising the above peptide and physiologically acceptable excipients for human and veterinary use. The preparation may comprise adjuvants for vaccination purposes. The administration route may be s.c., i.m., oral or nasal.



✓

decreasing the formation of A β protofibrils and/or lower meric forms thereof in a subject having, or suspected of having, AD.

The decreasing step above may be by active immunisation with a profibril or compound(s) with protofibril forming ability for prevention or treatment of Alzheimer's disease (AD), wherein said protofibril or compound(s) have enhanced protofibril forming ability and/or enhanced immunogenicity compared to the wild-type counterparts.

Alternatively, the decreasing step above is by passive immunisation with antibodies against protofibrils or compound(s) with protofibril forming ability, such as A β -Arc. The passive immunisation may be in combination with antibodies against other A β peptides with mutations/modifications leading to increased protofibril formation and/or immunogenicity, preferably AD related mutations.

Antibodies generated against the human A β sequence containing the Arctic mutation are directed towards A β protofibrils and therefore are of therapeutic value in the treatment of Alzheimer's disease. Because the A β peptide is in a protofibril conformation when used as an immunogen, antibodies against A β protofibrils are generated. Availability of such antibodies opens up possibilities for the development of an efficient and lasting vaccination for the prevention and treatment of Alzheimer's disease.

In another alternative the decreasing step of the method according to the invention is by administration of agents with anti-protofibrillar activity.

In yet a further aspect of the invention, a combination of the vaccine or passive immunization with monoclonal antibodies or compounds with anti-fibrillar activity with one or several other AD treatments such as, acetylcholinesterase inhibitors, nootropics, anti-inflammatory drugs, estrogen, neurotrophic factor agonists, β -secretase inhibitors, γ -secretase inhibitors and α -secretase agonists, can improve AD treatment efficacy. The rationale is that these substances/treatments work with completely different mechanisms of action and hence can be combined to the benefit for the AD patient.

Detailed description of the invention

The basis of the present invention is a pathogenic amyloid precursor protein (APP) mutation located within the A β sequence at codon 693 (E693G), causing AD in a family from northern Sweden. Surprisingly, carriers of this "Arctic" mutation show decreased A β 42 and A β 40 levels in plasma. This finding is corroborated *in vitro*, where the A β 42

concentration was low in conditioned media from cells transfected with APP_{E693G}. Fibrillization studies demonstrate that A β peptides with the Arctic mutation (A β 40Arc) form protofibrils at a much higher rate and in larger quantities than wild-type (wt) A β (A β 40wt). The unique finding of decreased A β plasma levels in the Arctic AD family highlights the complexity of the disease and is likely to reflect a novel pathogenic mechanism. The mechanism disclosed in the present invention involves a rapid A β protofibril formation leading to accelerated build-up of insoluble A β intra- and/or extracellularly.

In the present invention, the single amino acid substitution Glu to Gly at position 22 in the A β 4040Arc molecule was found to cause a dramatic increase in rate and capacity to form protofibrils compared to the A β 40wt peptide. Thus, when A β 42Arc and A β 40Arc are formed in the brain it is likely that they are more prone to be retained by cellular systems since the accelerated drive to form protofibrils enhances both A β -bulk and insolubility. Thus, factors promoting protofibril formation should be considered in the pathogenesis of sporadic AD. Increased protofibril formation is probably also operating in these more common forms of the disease. Indeed, the findings of the present invention open new avenues for possible therapeutic intervention using drugs targeted at preventing protofibril formation.

Studies on the Arctic mutation of the present invention have demonstrated a previously not described pathogenic mechanism for Alzheimer's disease through increased formation of A β protofibrils. A β with the Arctic mutation formed more stable protofibrils and at a much higher rate and in larger quantities than wild-type A β , even in the presence of equimolar amounts of wild-type A β . The formation is accelerated at least 2-10 times compared to protofibril formation of wild type A β peptides. The implication of this finding is that the dangerous species in the amyloid forming pathway that eventually leads to Alzheimer's disease is not the A β fibrils, but a form of the peptide that appears earlier in the fibril maturation process, the protofibrils. One implication of the findings related to the present invention is that it is important to prevent the formation of protofibrils in order to be able to prevent and treat Alzheimer's disease.

Non-human animals comprising transgenes which encode Arctic mutation APP can be used commercially to screen for agents having the effect of lowering the formation of A β protofibrils. Such agents can be developed as pharmaceuticals for treating abnormal APP processing and/or Alzheimer's disease, amongst other neurodegenerative conditions in

humans and animals, such as dogs. The transgenic animals of the present invention exhibit abnormal APP processing and expression, and can be used for pharmaceutical screening and as disease models for neurodegenerative diseases and APP biochemistry.

Figure legends

The present invention will now be further described with reference to the enclosed figures, in which:

Figure 1 shows kinetics of soluble forms of A β 1-40wt (a), A β 1-40Arc (b) and protofibril formation of A β 1-40wt, A β 1-40Arc vs a mixture of A β 1-40wt + Arc (1:1) (c). The A β 1-40Arc peptide (92 μ M) rapidly forms protofibrils (black dots) in comparison to the A β 1-40wt peptide (88 μ M), which mainly is in monomeric/dimeric (grey dots) form, data is taken from one experiment, representative of three (a and b). The protofibril formation rate was monitored during the first seven hours and the kinetics for the pure peptides (A β 1-40wt and A β 1-40Arc at 50 μ M) was compared to the protofibril formation rate of a 1:1 mixture (50 μ M) of A β 1-40wt + Arc (c).

Figure 2 depicts elution profiles showing A β 40wt (a-c) versus A β 40Arc (d-f) at 5 (a,d), 45 (b,e) and 125 (c,f) min of incubation. Accelerated protofibril (p) formation along with a parallel decline in the monomeric/dimeric (m/d) A β levels could be observed for A β 40Arc (d-f) as compared to A β 40wt (a-c). Data is from one experiment, representative of four. Initial peptide concentrations were 143 μ M and 138 μ M for A β 40wt and A β 40Arc, respectively.

EXAMPLES

The following examples are provided for illustration and are not intended to limit the invention to the specific example provided.

Example 1: Identification of the Arctic mutation

An APP mutation (E693G) in a family from northern Sweden, named the "Arctic" family, is identified, which spans over four generations. The family was screened for mutations in exons 16 and 17 of the APP gene by single strand conformation polymorphism analysis (SSCP) (L. Forsell, L. Lannfelt, (1995)). An abnormal mobility pattern was observed in

exon 17. Sequencing revealed an A→G nucleotide substitution, representing a glutamic acid to a glycine substitution at APP codon 693 (E693G), corresponding to position 22 in the Aβ sequence. Venous blood was drawn into tubes containing EDTA and DNA was prepared according to standard procedures. SSCP was performed. To sequence exon 17 of the APP gene a 319 bp fragment was amplified with the following primers 5'-CCT CAT CCA AAT GTC CCC GTC ATT-3' and 5'-GCC TAA TTC TCT CAT AGT CTT AAT TCC CAC-3'. The PCR products were purified with QIAquick PCR purification kit (Qiagen) prior to sequencing. Direct sequencing was performed in both 3' and 5' direction using the same primers and the BIG Dye cycle sequencing protocol (PE Biosystems) and were then analyzed on an ABI377 automated sequencer (PE Biosystems). The Arctic mutation was seen in one family and not in 56 controls or 254 cases with dementia. Carriers of the arctic mutation showed no vascular symptoms. The mutation was further verified by restriction analysis, since it destroyed a MbolI restriction site. The mutation was fully penetrant as no escapees were found. Two-point linkage analysis was performed between the mutation and affection status in the family with an age-dependent penetrance, giving a lod score of 3.66 at recombination fraction 0.00. Two-point lod score was calculated using Mlink from the linkage package (version 5.1) at each of the following recombination fractions 0.00, 0.10, 0.20, 0.30 and 0.40 (q males=q females). A single-locus model with an autosomal dominant inheritance was assumed, which was compatible with the inheritance as it appeared in the pedigree. A cumulative age dependent penetrance was assigned from the known ages of onset in the family. Individuals were put into different liability classes depending on the age at onset (affected) or age at last examination (unaffected). The disease gene frequency and the marker allele frequency were estimated to be 0.001 and the phenocopy rate was set to 0.0001.

Example 2: Clinical symptoms in carriers of the Arctic mutation

The family with the "Arctic" mutation was clinically and genealogically investigated. In this family, the mean age of onset was 56.6 years and the mean duration of the disease was 7 years (n=5).

The first symptom in most cases in this family was an insidious loss of memory for recently acquired information. Symptoms before clinical manifestation of Alzheimer's disease were decreased power of concentration and difficulties in handling stress situations. All affected individuals in generation IV had an early retirement pension because of the disease. The patients in generation IV were investigated by magnetic

resonance imaging (MRI), computed tomography (CT) and electroencephalography (EEG) which confirmed the diagnosis of Alzheimer's disease. In four individuals CT and MRI did not demonstrate signs of stroke or cerebral haemorrhage.

Example 3: Decreased A β plasma levels in carriers of the Arctic mutation

Pathogenic APP mutations have been shown to affect APP processing, as reflected in an increase of either total A β or A β 42 in the plasma of affected family members. The Arctic mutation is located in a region different from other AD-causing mutations. It was investigated as to whether the mutation manifested itself by affecting A β plasma levels. Plasma from nine mutation carriers, of which four were symptomatic, and eleven non-carriers in the family, were analysed by well-characterized sandwich ELISA systems, specifically detecting A β 42 (BAN50/BC05) and A β 40 (BAN50/BA27) (Suzuki et al. 1994)). To reassure that the Arctic mutation did not change any of the antibody recognition sites A β 40wt and A β 40Arc peptides were tested and found to be recognized equally well. Furthermore, plasma was spiked with synthetic peptides, revealing that both A β Arc and A β wt peptides were recovered by ELISA to the same extent. The data obtained was analyzed by non-parametric Mann-Whitney analysis. The A β 42 plasma concentration was 11.7 ± 3.9 fmol/ml and 16.0 ± 5.6 fmol/ml in mutation carriers and non-carriers, respectively, representing a 27% reduction of A β 42 in the mutation carriers ($p=0.04$). The A β 40 plasma concentration was 105 ± 22 fmol/ml and 141 ± 34 fmol/ml in mutation carriers and non-carriers, respectively, representing a 26% reduction of A β 40 in the mutation carriers ($p=0.01$). The A β 42/40 ratio was calculated for each individual, but no significant difference was found ($p=0.13$). In conclusion, concentrations of both A β 42 and A β 40 were unexpectedly and significantly reduced in individuals carrying the Arctic mutation.

Example 4: A β levels in cell culture

The effect of the Arctic mutation on A β formation was further investigated *in vitro* in transiently transfected HEK293 cells. APPwt was compared to the following mutations: Arctic (APP_{E693G}), Dutch (APP_{E693Q}), Italian (APP_{E693K}) and Flemish (APP_{A692G}). Constructs containing the Swedish double mutation (APP_{Swc}) and one APP mutation at codon 717 (APP_{V717F}), both with well-studied APP processing characteristics (Hardy (1997)), were used as positive controls. The mutations were introduced to APP695 cDNA in pcDNA3 using QuikChange™ Site-Directed Mutagenesis Kit according to the manufacturers

Sub
E4

instructions (Stratagene). The mutated constructs were verified by sequencing. For the ELISA measurements, HEK293 cells were seeded in six-well dishes and transfected with the different constructs using FUGENE™ 6 Transfection Reagent (Roche Diagnostics) according to the manufacturers instructions. 24 h after transfection, the cells were conditioned 48 h in OptiMEM containing 5% newborn calf serum. After withdrawal of the media for ELISA measurements, the APP expression in the cells were investigated by western blot using monoclonal antibody 22C11 (Roche Diagnostics). Media was conditioned and analyzed for A β levels by the same A β 42- and A β 40-specific sandwich ELISA systems as used for human plasma (Citron, et al. (1997)). The A β 42 and A β 40 concentrations and A β 42/40 ratios are shown in Table 1.

Table 1 A β 42/40 ratio and A β 42 and A β 40 levels in conditioned media from transiently transfected HEK293 cells

	A β 42/40 ratio	A β 42 \pm SD	A β 40 \pm SD
APP constructs	(%) \pm SD	(fmol/ml)	(fmol/ml)
APPwt	9.6 \pm 0.7	13.8 \pm 1.0	144 \pm 6
Arctic (E693G)	7.5 \pm 0.5*	11.2 \pm 0.6	149 \pm 3
Dutch (E693Q)	6.6 \pm 0.6*	9.6 \pm 0.7	147 \pm 12
Italian (E693K)	6.4 \pm 0.6*	8.0 \pm 0.7	126 \pm 17
Flemish (A692G)	11.7 \pm 1.6*	27.0 \pm 2.0	232 \pm 25
Mock (vector only)	7.2 \pm 2.4	2.1 \pm 1.0	28 \pm 5

* P=0.004 in comparison to APPwt

Decreasing A β 42/A β 40 ratios could be seen with all mutations at APP 693 (Arctic, Dutch, Italian). This may be due to increased rate of intracellular protofibril formation.

Example 5: Effect of Arctic mutation on protofibril formation

The effect of the single amino acid substitution (Glu22Gly) on amyloid fibrillization kinetics was investigated. Synthetic A β 1-40 was dissolved in physiological buffer and incubated for different periods of time. After centrifugation, the soluble A β in the supernatant, both low molecular weight (monomeric/dimeric) A β and protofibrils, were separated and analyzed using size exclusion chromatography (SEC) with UV detection at 214 nm. The morphology of the sedimented insoluble A β was visualized using negative stain and transmission electron microscopy (TEM).

$A\beta$ 1-40wt was purchased from Bachem, B ubendorf, Switzerland or Biosource International/QCB (Camarillo, CA, USA) and $A\beta$ 1-40Arc from Biosource International/QCB. The peptides were trifluoroacetic salts. They were stored at -20°C . All other chemicals were of highest purity available. Samples of each peptide were incubated, without agitation, at 30°C in 50 mM $\text{Na}_2\text{HPO}_4 \cdot \text{NaH}_2\text{PO}_4$ (pH 7.4) containing 0.1 M NaCl, for various time-points. Initial peptide concentrations were within the range of 88-143 μM , and were similar for both peptides in each experiment. After centrifugation (17 900 x g for 5 min at 16°C) monomeric/dimeric and protofibrillar $A\beta$ 1-40, sampled from the supernatant, were separated using SEC. A Merck Hitachi D-7000 LaChrom HPLC system, having a diod array detector model L-7455, a L-7200 model autosampler and a model L-7100 pump, coupled to a Superdex 75 PC3.2/30 column (Amersham Pharmacia Biotech, Uppsala, Sweden), was used for the chromatographic separation and analysis. Samples were eluted at a flow rate of 0.08 ml/min (ambient temperature) using 50 mM $\text{Na}_2\text{HPO}_4 \cdot \text{NaH}_2\text{PO}_4$ (pH 7.4), 0.15 M NaCl. Chromatograms were obtained by measuring UV absorbance at 214 nm. Peak areas for monomeric/dimeric and protofibrillar $A\beta$ were integrated using Merck-Hitachi Model D-7000 Chromatography Data Station Software. The mean of triplicate integrated peak values from the SEC measurements were used to generate each data point shown in Fig. 1 and 2. In addition, a standard curve was produced by correlating integrated peak areas with peptide concentrations as determined by quantitative amino acid analysis. The concentrations of total (at $t=0$ h) and soluble peptides remaining in solution after centrifugation were calculated from the standard curve.

SEC analysis of freshly dissolved $A\beta$ 1-40wt generated a single elution peak at a retention time of about 20 min (Fig. 2a). This peak represented the monomeric/dimeric forms of $A\beta$ 1-40wt (Walsh et al.(1997)). With increasing incubation time a second distinct peak appeared in the gel-excluded fraction with a retention time of about 12 min. This earlier peak contained protofibrils (Fig. 2b, c), as verified by ultracentrifugation, negative stain and TEM of $A\beta$ 1-40wt (data not shown), in line with previous findings (Walsh et al. (1997)). Similar retention times were obtained for the $A\beta$ 1-40Arc peptide (Fig. 2d-f). However, $A\beta$ 40Arc generated protofibrils much faster and in larger quantities than $A\beta$ 40wt. Chromatograms from three early time-points of incubation illustrate this difference (Fig. 1). The monomeric/dimeric $A\beta$ 40Arc peak declined in parallel with the growth of the

protofibrillar peak (Fig. 2d-f). The maximum concentration (111 μM) of A β 40Arc protofibrils was observed at 6.5 h.

Kinetic studies up to 48 h showed that A β 1-40wt generated a small quantity of protofibrils with a maximum concentration at 25 h (Fig. 1a). In contrast, a rapid and significant formation of protofibrils was seen within the first 5 h of incubation with a simultaneous rapid decline in the concentration of the monomeric/dimeric A β 1-40Arc peptide (Fig. 1b). Since carriers of the Arctic mutation are heterozygotes they generate both A β wt and A β Arc. Assuming equimolar in vivo production, the kinetics of protofibril formation was studied in a 1:1 mixture of A β 1-40wt and A β 1-40Arc. This mixture of peptides showed kinetics that were intermediate to the single peptide curves (Fig. 1c).

Example 6: Morphology of A β -Arc

A typical fibrillar morphology of A β 1-40Arc in sedimented samples from kinetic studies was confirmed by negative stain and TEM. A β peptide samples were prepared and incubated as indicated for the kinetic studies, using higher peptide concentrations (617 μM). After 8 days, aggregated A β species were sedimented using the same centrifugation parameters as described above. Buffer was removed and pelleted material was suspended in 50 μl water using gentle sonication (2 x 6s). Eight μl samples were applied to carbon stabilized Formvar film grids (Ted Pella, Inc., Redding, CA, USA). Samples were negatively stained with 8 μl uranyl acetate (1%) (E. Merck, Darmstadt, Germany). Four grids were prepared for each sample and examined using a Philips CM10 TEM. Samples from pellets sedimented during the kinetic experiments were also examined. Similar to the sedimented A β 40wt, large mesh-works of A β could be seen in these preparations. Protofibrils could also be discerned in the sedimented samples. The A β 1-40Arc protofibrils were longer and less curved compared to the A β 1-40wt protofibrils. Inter-twining of several fibrils was more common in the A β 40Arc preparations, resulting in larger fibril diameters.

Example 7: Kinetic studies

Kinetic studies comparing the formation of A β 40gly22 protofibrils in the presence of a high and a low concentration of NaCl:

The experiments examining A β 40gly22 protofibril and fibril formation, have been performed in 50 mM phosphate buffer supplemented with 100 mM NaCl. They present

References

- Citron, et al. *Nature Med* 3, 67-72 (1997).
- Conway, et al., *Proc Natl Acad Sci USA* 97, 571-576 (2000).
- De Jonghe, et al., *Neurobiol Disease* 5, 281-286 (1998).
- Forsell, Lannfelt, *Neurosci Lett* 184, 90-93 (1995).
- Giulian et al., *J Biol Chem*, 273, 29719-19726, (1998).
- Grabowski et al., *Ann Neurol* 49, 697-705 (2001)
- Hardy, *Trends Neurosci.* 20, 154-159 (1997).
- Harper et al., *Biochemistry* 38, 972-8980 (1999).
- Hendriks, et al., *Nature Genet* 1, 218-221 (1992).
- Kamino, et al., *Am J Hum Genet* 51, 998-1014 (1992).
- Levy, et al., *Science* 248, 1124-1126 (1990).
- Mullan, et al., *Nature Genet* 1, 345-347 (1992).
- Palmert et al. *PNAS* 86:6338 (1989)
- Schenk et al., *Nature*, 400, 173-177 (1999)
- Scheuner, et al., *Nature Med* 2, 864-869 (1996).
- Serpell L.C. *Biochim. Biophys. Acta*, 1502, 16-30 (2000).
- Sherrington et al. *Nature* 375:754 (1995)

Suzuki, et al., *Science* **264**, 1336-1340 (1994).

Tagliavini, et al., *Alz Report* 2, S28 (1999).

Walsh et al., *J Biol Chem* **36**, 25945-25952 (1999).

Weidemann et al. *Cell* 57:115 (1989)

Wirak et al. *Science* 253:323 (1991)

SEQUENCE LISTING

<110> Lannfelt, Lars

<120> Prevention and treatment of Alzheimer's disease

<130> P05384PC

<210> 1

<211> 42

<212> PRT

<213> Homo sapiens

<400> 1

Asp	Ala	Glu	Phe	Arg	His	Asp	Ser	Gly	Tyr	Glu	Val	His	His	Gln	Lys	Leu
1				5				10						15		

Val	Phe	Phe	Ala	Gly	Asp	Val	Gly	Ser	Asn	Lys	Gly	Ala	Ile	Ile	Gly	Leu
			20				25						30			

Met	Val	Gly	Gly	Val	Val	Ile	Ala
35						40	

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2